

# HASH BIOTECH LIMITED

A GROUP COMPANY OF HASH GROUP



## C-PHYCOCYANIN

### CHARACTERISTICS

MOLECULAR WEIGHT	: 232KD <sub>a</sub>
COMPOSITION	: C-PHYCOCYANIN COMPRISES OF TWO SUBUNITS $\alpha$ AND $\beta$ OF MOLECULAR WEIGHT IN THE RANGE OF 18KD <sub>a</sub> & 20KD <sub>a</sub> RESPECTIVELY.
PURITY	: A <sub>620</sub> /A <sub>280</sub> 3.5-4
ABSORPTION MAXIMA	: 620 nm
EMISSION MAXIMA	: 647 nm
EXTINCTION COEFFICIENT	: E <sub>1%</sub> AT 620 = 70
ISOELECTRIC POINT	: 4.65
SOURCE	: <i>Spirulina platensis</i>
ORIGIN	: INDIA


C-PHYCOCYANIN (C-PC) OCCURS AS THE CYANOBACTERIAL PHYCOBILIPROTEIN WITH HIGH FLUORESCENCE QUANTUM YIELD, MOLAR EXTINCTION COEFFICIENTS, ABSORPTIVITY AND STABILITY. IT CAN BE EASILY CROSS LINKED WITH ANTIBODIES AND OTHER PROTEINS BY USING CONVENTIONAL MOLECULAR TAGGING TECHNIQUES WITHOUT LOSING ITS FLUORESCENT PROPERTIES.

BECAUSE OF ITS MIRACULOUS FLUORESCENCE PROPERTIES, IT MAY BE USED AS A FLUORESCENT BIOMARKER WITH PROTEINS, ANTIBODIES AND NUCLEIC ACIDS FOR DIAGNOSING THE VARIOUS LETHAL DISEASES AND IN DIFFERENT DIAGNOSTIC KITS. IT IS ALSO USED AS A STRAIGHT MARKER IN VARIOUS APPLICATIONS LIKE FRET ASSAYS, FLUORESCENCE MICROSCOPY, FLUORESCENCE ACTIVATED CELL SORTING (FACS), FLUORESCENCE IN SITU HYBRIDIZATION (FISH), FLUORESCENCE CORRELATION SPECTROSCOPY (FCS), GEL ELECTROPHORESIS, ISOELECTRIC FOCUSING AND GEL EXCLUSION CHROMATOGRAPHY.

THE FILAMENTOUS PROKARYOTIC CYANOBACTERIUM *Spirulina platensis* IS A RICH SOURCE OF PHYCOCYANIN WHICH ALSO POSSESSES A WIDE RANGE OF COLORED COMPONENTS I.E. CHLOROPHYLL, ALLOPHYCOCYANIN (APC), PHYCOERYTHRIN (PE) AND CAROTENOIDS. [1][5].

THE PHYCOBILIPROTEINS ARE ANTENNAE-PROTEIN PIGMENTS INVOLVED IN LIGHT HARVESTING AND ARE ORGANIZED IN SUPRAMOLECULAR COMPLEXES, CALLED PHYCOBILISOMES (PBSS), THOSE ARE ASSEMBLED IN REGULAR ARRAYS ON THE OUTER SURFACE OF THE THYLAKOID MEMBRANES.

IN CYANOBACTERIA FOUR MAIN CLASSES OF PHYCOBILIPROTEINS EXIST: ALLOPHYCOCYANIN (APC, BLuish GREEN), PHYCOCYANIN (PC, BLUE), PHYCOERYTHRIN (PE, PURPLE), AND PHYCOERYTHROCYANIN (PEC, ORANGE).




MOLECULAR SKELETON OF C-PHYCOCYANIN

PHYCOBILIPROTEINS ARE OLIGOMERIC PROTEINS, BUILT UP FROM CHROMOPHORE-BEARING POLYPEPTIDES BELONGING TO TWO FAMILIES ( $\alpha$  AND  $\beta$ ) PROBABLY ORIGINATING FROM A COMMON ANCESTOR [2].


THE COLORS OF PHYCOBILIPROTEINS ORIGINATE MAINLY FROM COVALENTLY BOUND PROSTHETIC GROUPS THAT ARE OPEN-CHAIN TETRAPYRROLE CHROMOPHORES BEARING A, B, C AND D RINGS NAMED PHYCOBILINS.


THESE CHROMOPHORES ARE GENERALLY BOUND TO THE POLYPEPTIDE CHAIN AT CONSERVED POSITIONS EITHER BY ONE CYSTEINYL THIOESTER LINKAGE THROUGH THE VINYL SUBSTITUENT ON THE PYRROLE RING A OF THE TETRAPYRROLE OR OCCASIONALLY BY TWO CYSTEINYL THIOESTER LINKAGES



THREE DIMENSIONAL MOLECULAR STRUCTURE OF C-PHYCOCYANIN

- OXYGEN
- CARBON
- NITROGEN
- HYDROGEN





SPECTROSCOPIC CHARACTERIZATION OF PURIFIED C-PC IN 5 mM SODIUM PHOSPHATE BUFFER, PH 7.0. ABSORPTION SPECTRUM OF C-PC, SPECTRUM RECORDED AT ROOM TEMPERATURE.

TABLE SHOWING SPECTROSCOPIC CHARACTERS OF PIGMENTS[3]

S.No.	PIGMENT	ABSORPTION MAXIMA(IN nm)	EMISSION MAXIMA(IN nm)
1.	ALLO-PHYCOCYANIN	650-655	660
2.	C-PHYCOCYANIN	615-640	637
3.	PHYCOERYTHRIN	565-575	577

#### APPLICATION OF C-PHYCOCYANIN

PHYCOBILIPROTEINS PLAY AN IMPORTANT ROLE IN FLUORESCENT BASED DETECTION SYSTEMS, PARTICULARLY FOR FLOW CYTOMETRY. THE SPECTRAL PROPERTIES, SUCH AS:

- EXCITATION AND EMISSION AT THE RED END OF THE SPECTRUM, WHERE INTERFERENCE FROM BIOLOGICAL MATRICES TEND TO BE LESS.
- A LARGE STOKES SHIFT, SO THAT INTERFERENCE FROM RAYLEIGH AND RAMAN SCATTER AND OTHER FLUORESCING COMPONENTS IS LESS SIGNIFICANT OR NONEXISTENT.
- IMMUNITY FROM QUENCHING BY NATURALLY OCCURRING BIOLOGICAL SUBSTANCES.
- HIGH SOLUBILITY IN AQUEOUS ENVIRONMENT SO THAT NONSPECIFIC BINDING EFFECTS ARE MINIMAL.

SYNTHESIS OF CONJUGATES OF PHYCOBILIPROTEINS WITH MOLECULES HAVING BIOLOGICAL SPECIFICITY, LIKE IMMUNOGLOBULINS, PROTEIN A, BIOTIN AND AVIDIN, WERE REPORTED AND SHOWED THAT PHYCOBILIPROTEINS CONJUGATES ARE EXCELLENT REAGENTS FOR TWO COLOR FLUORESCENCE ANALYSIS OF SINGLE CELLS USING FLUORESCENCE ACTIVATED CELL SORTER (FACS) [4].

A PHOTOCHEMICAL METHOD IS DESCRIBED FOR TREATING CANCER WHEREIN PHYCOCYANIN IS ADMINISTERED TO A PATIENT SUFFERING FROM INTERNAL OR SKIN CANCER. ONCE ADMINISTERED, PHYCOCYANIN IS SELECTIVELY TAKEN UP INTO CANCER CELLS, AND UPON SUBSEQUENT IRRADIATION DESTRUCTION OF THE CANCER CELLS OCCURS.

THE PHARMACOLOGICAL PROPERTY ATTRIBUTED BY PHYCOCYANIN INCLUDES ANTIOXIDANT, ANTI-INFLAMMATORY, EUROPROTECTIVE AND HEPATOPROTECTIVE ACTIVITY.

WHEN IT WAS EVALUATED AS AN ANTIOXIDANT *in vitro*, IT WAS ABLE TO SCAVENGE ALKOXYL, HYDROXYL AND PEROXYL RADICALS, AND INHIBITS MICROSO MAL LIPID PEROXIDATION INDUCED BY  $Fe^{+2}$ -ASCORBIC ACID OR THE FREE RADICALS INITIATORS 2, 2' AZOBIS (2- AMIDINOPROPANE) DIHYDROCHLORIDE, (AAPH). THEY ALSO REDUCED EDEMA, HISTAMINE (HI) RELEASE, MYELOPEROXIDE (MPO) ACTIVITY AND THE LEVELS OF PROSTAGLANDIN (PGE2) AND LEUKOTRIENE (LTB4) IN THE INFLAMED TISSUES [6][7].

PHYCOCYANIN SELECTIVELY IMPREGNATE BIOLOGICAL STRUCTURES INCLUDING ATHEROSCLEROTIC PLAQUE AND TUMORS. IMAGING OF BIOLOGICAL STRUCTURES CAN BE ENHANCED BY TAGGING TO A PHYCOCYANIN AN ATOM WHICH IS COMPATIBLE WITH THE COMPLEMENTARY TO ANY OF THE IMAGING MODALITIES [9].

THE TAGGED PHYCOCYANIN CAN THEN BE ADMINISTERED TO A SUBJECT. THE IMPROVEMENT CAN BE UTILIZED TO ENHANCE IMAGES OF ALL STRUCTURES, ARTERIAL WALL THICKNESS, ATHEROSCLEROTIC PLAQUE, LUMINAL BOUNDARIES AND TO BETTER DELINEATE TUMORS MASS OUTLINES [8].

C-PHYCOCYANIN DERIVED FROM *Spirulina platensis* POWERFULLY INFLUENCED SERUM CHOLESTEROL CONCENTRATIONS AND IMPARTED A STRONGER HYPOCHOLESTEROLEMIC ACTIVITY[10]

#### REFERENCES:

- [1]HENRIKSON R (1989), IN: HENRIKSON ROBERT, EDITOR. EARTH FOOD *Spirulina*. LAGUNA BEACH, CA: RONORE ENTERPRISES INC.
- [2]GLAZER AN (1989), LIGHT GUIDES - DIRECTIONAL ENERGY TRANSFER IN A PHOTOSYNTHETIC ANTENNA. J CELL BIOL 264(5):1-4.
- [3]BRYANT DA, GUGLIELMI G, TANDEAU DE MARSAC N ET AL (1979) THE STRUCTURE OF CYNOBACTERIAL PHYCOBILISOMES: A MODEL. ARCH MICROBIOL 123:113-127
- [4]GLAZER AN (1985), LIGHT HARVESTING BY PHYCOBILISOMES. ANNU REV BIOPHYS CHEM 14:47-77.
- [5]DI VT, GLAZER AN, STRYER L (1982), FLUORESCENT PHYCOBILIPROTEIN CONJUGATES FOR ANALYSIS OF CELLS AND MOLECULES. J CELL BIOL 93:981-986.
- [6]SAMY BOUSSIBA AND AMOS E. RICHMOND(1979), ISOLATION AND CHARACTERIZATION OF PHYCOCYANINS FROM THE BLUE-GREEN ALGA *Spirulina platensis*.ARCH. MICROBIOL. 120, 155-159.
- [7]BHAT VB, MADYASTHA KM (2000), C-PHYCOCYANIN : A POTENT PEROXYL RADIAL SCAVENGER IN VIVO AND IN VITRO. BIOCHEM BIOPHYS RES COMM 275(1):20-25.
- [8]ROMAY CH, GONZALEZ R, LEDON N ET AL (2003), C-PHYCOCYANIN A BILIPROTEIN WITH ANTIOXIDANT, ANTI-INFLAMMATORY AND NEUROPROTECTIVE EFFECTS. CURRENT PROTEIN PEPTIDE Sci 4:207-216.
- [9]NABIL CM (1998), ANATOMICAL IMAGING WITH PHYCOCYANINS. US PATENT 5807536, 15 SEP 1998.
- [10]NAGAOKA S, SHIMIZU K, KANEKO H ET AL (2005) A NOVEL PROTEIN C-PHYCOCYANIN PLAYS A CRUCIAL ROLE IN THE HYPOCHOLESTEROLEMIC ACTION OF *Spirulina platensis* CONCENTRATE IN RATS. J NUTR 135:2425-2430

